

Anomalous single production of the fourth generation charged leptons at future ep colliders

A. K. Çiftçi^{1,*}, R. Çiftçi², H. Duran Yıldız^{3,†}, S. Sultansoy^{4,5}

¹⁾ Physics Department, Faculty of Sciences, Ankara University, 06100 Tandogan, Ankara, Turkey,

²⁾ A. Yüksel Cad. 7/15, 06010 Etlik, Ankara, Turkey,

³⁾ Physics Department, Faculty of Sciences and Arts, Dumlupınar University, Merkez Kampüs, Kütahya, Turkey,

⁴⁾ Physics Department, Faculty of Sciences and Arts, Gazi University, 06500 Teknikokullar, Ankara, Turkey,

⁵⁾ Institute of Physics, Academy of Sciences, H. Cavid Avenue 33, Baku, Azerbaijan

Corresponding authors: ^{*}ciftci@science.ankara.edu.tr, [†]hyildiz@dumlupinar.edu.tr

Abstract

Possible single productions of the fourth standard model family charged leptons via anomalous interactions at the future ep colliders are studied. Signatures of such anomalous processes and backgrounds are discussed in detail.

PACS: 12.60.-i, 14.60.-z, 14.80.-j.

1. Introduction

A new era in High-Energy Physics will be opened with the construction of TeV scale colliders [1] in coming years. The first of the such colliders is the Large Hadron Collider (LHC) with 14 TeV center of mass energy and $10^{34} \text{ cm}^{-2}\text{s}^{-1}$ design luminosity. As for the lepton colliders, there are two realistic proposals namely International Linear Collider (ILC) with 0.5 TeV center of mass energy and $2 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$ luminosity [2] and Compact Linear Collider (CLIC) with 1 TeV (second stage) center of mass energy and $2.7 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$ luminosity [3]. Construction of ILC or CLIC tangential to the LHC ring will give opportunity to build multi-TeV center of mass energy lepton-hadron colliders.

The latest lepton-hadron collider was HERA with $\sqrt{s} \approx 0.3 \text{ TeV}$. One could build the next lepton-hadron collider by adding an electron ring at LHC tunnel [4]. In our opinion even though it is feasible, to build linac-ring type ep collider is better because of its versatility (possibility of building γp , γ -nucleus and FEL γ -nucleus colliders). Depending on its center of mass energy lepton-hadron colliders (ILC or CLIC on LHC) are named as QCD-Explorer with $\sqrt{s} = 1.4 \text{ TeV}$ [5, 6] and Energy Frontier-ep collider with $\sqrt{s} = 3.74 \text{ TeV}$ [7] (1 fb^{-1} integrated luminosity for both options). QCD-Explorer is a necessary machine for adequate interpretation of LHC data. Differing from ring-ring type machine, one could increase the center of mass energy of the collider by increasing the length of the linac (adding more units of ILC or CLIC accelerating structures). Eventually, it is possible to reach frontier energies by using full size of the ILC or CLIC. TeV scale hadron, lepton and lepton-hadron colliders will be complementary each other from the view point of physics at TeV scale [1, 5]. For this reason different new phenomena should be studied comparatively in these colliders. LHC, ILC and CLIC are well developed with respect to physics potential. However, there is little study on search potential of ep colliders. Therefore, they deserve further attention.

It is known that flavor democracy hypothesis forces the existence of the fourth standard model (SM) family [8-10]. The masses of the fourth family fermions are expected to be nearly degenerate and lie between 300 and 700 GeV [11]. Due to large mass value, one could expect a serious

contribution of anomalous interactions for the fourth SM family. This paper is devoted to detailed study the anomalous single production of the fourth family charged leptons at above ep colliders.

2. Anomalous Interactions of the Fourth SM Family Lepton

The effective lagrangian for the anomalous interactions of the fourth family charged lepton can be adopted from [12] with minor modifications as:

$$L = \left(\frac{K_\gamma^{l_i}}{\Lambda} \right) e_l g_e \bar{l}_4 \sigma_{\mu\nu} l_i F^{\mu\nu} + \left(\frac{K_Z^{l_i}}{2\Lambda} \right) g_Z \bar{l}_4 \sigma_{\mu\nu} l_i Z^{\mu\nu} + h.c. , \quad (i = 1, 2, 3) \quad (1)$$

where $K_{\gamma,Z}^{l_i}$ are the anomalous couplings for the neutral currents with a photon and a Z boson, respectively. Λ is the cutoff scale for the new physics and e_l is the lepton charge; g_e and g_Z are the electroweak coupling constants, $g_Z = g_e / \cos\theta_W \sin\theta_W$, where θ_W is the Weinberg angle. In the Eq. (1) $\sigma_{\mu\nu} = i(\gamma_\mu \gamma_\nu - \gamma_\nu \gamma_\mu)/2$. $F^{\mu\nu}$ and $Z^{\mu\nu}$ are field strength tensors of the photon and Z boson, respectively. Obviously new interactions will lead to additional decay channels of the fourth family leptons. In order to calculate decay widths, we have implemented the new interaction vertices into the CompHEP [13]. Results of the calculations for l_4 assuming $(K/\Lambda) = 1 \text{ TeV}^{-1}$ are given in Table 1. Let us remind that decay widths are proportional to $(K/\Lambda)^2$.

At the same table, we present the widths of the l_4 SM decay modes, as well. While calculating these values, we have used leptonic CKM Matrix elements from [14], namely: $V_{\nu_4\tau} = 2.34 \times 10^{-5}$, $V_{\nu_4\mu} = 6.81 \times 10^{-4}$, $V_{\nu_4e} = 4.64 \times 10^{-4}$; $V_{l_4\nu\tau} = 8.14 \times 10^{-4}$, $V_{l_4\nu\mu} = 1.28 \times 10^{-4}$, $V_{l_4\nu e} = 6.43 \times 10^{-6}$. Let us remind that SM decay widths are proportional to $|V_{l_4\nu}|^2$. Depending on relative magnitudes of (K/Λ) and $|V_{l_4\nu}|$, SM or anomalous decays will dominate. The total decay width Γ of the fourth family leptons and the relative branching ratios are given in Fig. 1.

Table 1. Branching ratios (%) and total decay widths for $m_{l_4}(\text{GeV})$

m_{l_4}	$\gamma e(\mu, \tau)$	$Z\tau(e, \mu)$	$W\nu_\tau(\times 10^{-4})$	$W\nu_\mu(\times 10^{-5})$	$W\nu_e(\times 10^{-8})$	$\Gamma_{Tot}(\text{GeV})$
100	32	1.4(1.8)	5.3	1.3	2.9	0.01
150	20	13	9.1	2.2	4.9	0.06
200	17	16	8.8	2.2	4.8	0.18
300	15	18	8.3	2.1	4.5	0.70
700	14	19	7.9	1.9	4.3	9.54

3. Anomalous Single Production of the Fourth SM Family Leptons at ep Colliders

In ep colliders ($\sqrt{s} = 1.4 \text{ TeV}$ and $\sqrt{s} = 3.74 \text{ TeV}$ are considered), we investigate single anomalous production of the fourth SM family leptons. The calculated cross-sections for l_4 are displayed in Fig. 2 and Fig. 3. We consider $ep \rightarrow l_4 X \rightarrow \gamma \mu X$ and $ep \rightarrow l_4 X \rightarrow ZeX$ processes. In the former process anomalous decay of l_4 can be detected easily at ep colliders due to no background. Number of events for $m_{l_4} = 300$ (700) GeV at $\sqrt{s} = 1.4 \text{ TeV}$ and $\sqrt{s} = 3.74 \text{ TeV}$ with 1 fb^{-1} integrated luminosity are 240 (30) and 1170 (460), respectively.

However, the latter process has SM backgrounds. In order to extract the fourth generation lepton signal and to suppress the background, we impose cuts on the eZ invariant mass. Thus, in calculations, we used $|m_{eZ} - m_{l_4}| < 25 \text{ GeV}$ for the mass range $m_{l_4} = 100 - 1000 \text{ GeV}$ and $|m_{eZ} - m_{l_4}| < 50 \text{ GeV}$ for the mass range of $1 - 2.6 \text{ TeV}$ with $p_T^q > 10 \text{ GeV}$ cuts. All calculations for the signal and background were done by using the high-energy package CompHEP with CTEQ6L [15] distribution function. We present signal and background cross-sections with respect to invariant mass of eZ in Tables 2 and 3 for $\sqrt{s} = 1.4 \text{ TeV}$, $\sqrt{s} = 3.74 \text{ TeV}$ options, respectively.




Figure 1: (a) The total decay width Γ in GeV of the fourth family lepton and (b) the branching ratios (%) depending on the mass of the fourth family lepton.




Figure 2: The total production cross-section of the process $ep \rightarrow l_4 X$ as a function of m_{l_4} with the center of mass energy $\sqrt{s} = 1.4$ TeV.

In order to examine the potential of the collider to search for the investigation of the anomalous interactions of the fourth SM family lepton in detail, we define statistical significance (SS) of the signal:

$$SS = \frac{|\sigma_{S+B} - \sigma_B|}{\sqrt{\sigma_B}} \sqrt{L_{int}} \quad (2)$$

where L_{int} is the integrated luminosity of the collider. The values SS evaluated at each fourth SM family lepton mass points are shown in the last column of Table 2 and Table 3. As seen from tables, one obtains good SS for the expected mass range of leptons.

Table 2. Total cross section of signal and background for $\sqrt{s} = 1.4$ TeV and $(K/\Lambda) = 1\text{TeV}^{-1}$

$m_{l_4}(\text{GeV})$	$\sigma_{S+B}(\text{pb})$	$\sigma_B(\text{pb})$	SS
100	0.13	0.1	3
200	0.49	0.09	42.16
300	0.33	0.05	39.60
400	0.21	0.03	32.86
500	0.13	0.02	24.59
600	0.075	0.015	15.49
700	0.039	0.009	10.00
800	0.019	0.005	6.26
900	0.0076	0.002	3.96
1000	0.0028	0.0011	1.62

Fig. 4 shows the invariant mass m_{eZ} distribution in the reaction $ep \rightarrow l_4 X \rightarrow ZeX$ for the SM background and the signal with the inclusion of the fourth family lepton with masses $m_{l_4} = 300$ GeV, $m_{l_4} = 500$ GeV, and $m_{l_4} = 700$ GeV for QCD Explorer.




Figure 3: The total production cross-section of the process $ep \rightarrow l_4 X$ as a function of m_{l_4} with the center of mass energy $\sqrt{s} = 3.74$ TeV.

Fig. 5 shows the invariant mass m_{eZ} distribution in the reaction $ep \rightarrow l_4 X \rightarrow ZeX$ for the SM background and the signal with the inclusion of the fourth family lepton with masses $m_{l_4} = 500$ GeV, $m_{l_4} = 1$ TeV, and $m_{l_4} = 1.5$ TeV for Energy Frontier-ep collider.

Table 3. Total cross section of signal and background for
for $\sqrt{s} = 3.74$ TeV, $(K/\Lambda) = 1\text{TeV}^{-1}$

$m_{l_4}(\text{GeV})$	$\sigma_{S+B}(\text{pb})$	$\sigma_B(\text{pb})$	SS
200	1.234	0.064	146.25
400	0.856	0.026	162.77
600	0.554	0.014	144.32
800	0.339	0.009	110.00
1000	0.32	0.03	52.94
1200	0.19	0.02	38.01
1400	0.105	0.015	23.24
1600	0.062	0.012	14.43
1800	0.028	0.008	7.07
2000	0.016	0.006	4.08
2200	0.009	0.004	2.5
2400	0.004	0.002	1.41
2600	0.002	0.001	1

4. Conclusion

As a result of study it is seen that ep colliders are advantageous to investigate anomalous interactions of the fourth SM family charged lepton. QCD Explorer will cover fourth family charged lepton up to 900 GeV while at Energy Frontier-ep collider the masses up to 2100 GeV can be reached. The regions are similar to reaches for anomalous production of the fourth family neutrino [16]. Hadron colliders including LHC are not very good place to investigate fourth family leptons [17] whereas, the capacity of lepton colliders are limited by their center of mass energy.




Figure 4: Invariant mass m_{eZ} distribution of signal and background at QCD Explorer.




Figure 5: Invariant mass m_{eZ} distribution of signal and background at Energy Frontier-ep collider.

Acknowledgments

This work is supported by TAEK and DPT with grant number DPT2006K-120470. One of the authors (H. Duran Yıldız) supported by TÜBİTAK with the project number 105T442.

References

- [1] S. Sultansoy, Tr. J. of Phys. 22 (1998) 575-594.
- [2] B. Barish, ILC NewsLine, 7 December 2006.
- [3] E. Accomando et al., hep-ph/0412251.
- [4] J. B. Dainton et al., hep-ex/0603016.
- [5] S. Sultansoy, Eur. Phys. J. C 33 s01 (2004) 1064; Proc. of Knoxville 2005, Particle Accelerator Conference, pg 4329.
- [6] D. Schulte and F. Zimmermann, EPAC'04, Lucerne, Switzerland, p. 632; CERN-AB-2004-079, CLIC-NOTE-608.
- [7] F. Tecker (ed.), CLIC-Note-613.
- [8] A. Datta, S. Raychaudhuri, Phys. Rev. D 49 (1994) 4762.
- [9] A. Çelikel, A. K. Çiftçi, S. Sultansoy, Phys. Lett. B 342 (1995) 257.
- [10] S. Sultansoy, hep-ph/0004271; hep-ph/0610279.
- [11] S. Atağ et al., Phys. Rev. D 54 (1996) 5745-5749.
- [12] E. Arık et al., Phys. Rev. D 67 (2003) 035002.
- [13] A. Pukhov et al., hep-ph/9908288.
- [14] A. K. Çiftçi et al., Phys. Rev. D 72 (2005) 053006.
- [15] H. L. Lai et al., Phys. Rev. D 51 (1995) 4763.
- [16] A. Alan, A. T. Alan, Physics Letters B 632 (2006) 357-359.
- [17] ATLAS Physics TDR, Vol. II, 1999.